

MAIN PANEL

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**Tribute<sup>®</sup>**

**SELECTIVE TURF HERBICIDE**

**ACTIVE CONSTITUENT: 22.5 g/L FORAMSULFURON**

**GROUP B HERBICIDE**

**1 L**

For post-emergent control of certain grass weeds in turf as specified in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE**

## REAR PANEL

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**SAFETY DIRECTIONS**

May irritate the eyes. Will irritate the skin. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. When opening the container, preparing and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length chemical-resistant gloves. Wash hands after use.

**FIRST AID:** If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which may be found at [www.bayeres.com.au](http://www.bayeres.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 63240 / 45114

Tribute<sup>®</sup> and Destiny<sup>®</sup> are Registered Trademarks of Bayer.

BAR  
CODE

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

Bayer Environmental Science  
A Business Operation of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd, Hawthorn East, Vic, 3123  
Phone: (03) 9248 6888  
Technical Enquiries: 1800 804 479  
Website: [www.bayeres.com.au](http://www.bayeres.com.au)

Batch Number:

Date of Manufacture:

(label code)

**CAUTION  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**TRIBUTE SELECTIVE TURF HERBICIDE**

**ACTIVE CONSTITUENT:** 22.5 g/L FORAMSULFURON

<b>GROUP</b>	<b>B</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**For post-emergent control of certain grass weeds in turf as specified in the Directions for Use table.**

**READ THIS BOOKLET BEFORE USE**

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**SAFETY DIRECTIONS**

May irritate the eyes. Will irritate the skin. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. When opening the container, preparing and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length chemical-resistant gloves. Wash hands after use.

**FIRST AID:** If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which may be found at [www.bayeres.com.au](http://www.bayeres.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 63240 / 45114

**DIRECTIONS FOR USE****RESTRAINTS**

**DO NOT** apply with aircraft.

**DO NOT** apply through any type of irrigation equipment.

**DO NOT** apply with a nozzle height greater than 50cm above the ground.

**DO NOT** apply to turf or weeds under stress.

**DO NOT** apply on turf exhibiting injury from previous applications of other products.

**DO NOT** apply to turf which is not well established.

**DO NOT** apply to short mown turf (less than 10mm).

**DO NOT** apply if heavy rain has been forecast within 48 hours.

**DO NOT** apply to waterlogged soil.

**DO NOT** apply within two weeks of couchgrass sprigging.

**DO NOT** apply within four weeks prior to overseeding couchgrass with a cool season grass such as ryegrass, as some temporary stunting may occur for up to two weeks.

**SPRAY DRIFT RESTRAINTS**

**DO NOT** apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturers specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

**DO NOT** apply when wind speed is less than 3 or more than 20 km per hour as measured at the application site

**DO NOT** apply during surface temperature inversion conditions at the application site

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are: **1** date with start and finish times of application; **2** location address and paddock/s sprayed; **3** full name of this product; **4** amount of product used per hectare and number of hectares applied to; **5** crop/situation and weed/pest; **6** wind speed and direction during application; **7** air temperature and relative humidity during application; **8** nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; **9** name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

**MANDATORY NO-SPRAY ZONES**

**DO NOT** apply if there are aquatic or wetland areas including aquacultural ponds, surface streams and rivers within 80 metres downwind from the application area.

**DO NOT** apply if there are sensitive crops, gardens, landscaping vegetation, protected native vegetation or protected animal habitat within 30 metres downwind from the application area.

**DIRECTIONS FOR USE TABLE**

<b>SITUATION</b>	<b>WEEDS CONTROLLED</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Turf - only apply to couchgrass (common and hybrid couchgrass NOT Queensland Blue Couch)	Winter grass ( <i>Poa annua</i> ) Ryegrass ( <i>Lolium perenne</i> )	1.5 L/ha	A repeat application may be required if a dense weed cover or larger plants are present. Ryegrass control is improved with the addition of DESTINY at 25 g/ha.
	Crowsfoot Grass ( <i>Eleusine indica</i> )	2 L/ha	Two to three applications at 14-day intervals may be required for effective control.
	Winter grass ( <i>Poa annua</i> ) Ryegrass ( <i>Lolium perenne</i> ) Crowsfoot Grass ( <i>Eleusine indica</i> )	8 to 15 mL / L water	<u>Spot treatment only:</u> Use the higher rate for the more difficult to control or larger weeds (crowsfoot and larger ryegrass plants). Spray to wet, but do not apply beyond runoff.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**DO NOT GRAZE TREATED TURF OR FEED TURF CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK**

**DO NOT USE CLIPPINGS FROM TREATED AREAS FOR MULCH AROUND VEGETABLES OR FRUIT TREES**

**GENERAL INSTRUCTIONS**

TRIBUTE Selective Turf Herbicide is a selective sulfonylurea herbicide. It is predominantly a foliar herbicide with less activity via the soil. TRIBUTE will not reliably control weeds that emerge after spraying. Best results are achieved when the product is applied under good growing conditions and to weeds and turf which are actively growing and not under stress. Avoid mowing during the three to four days preceding or following treatment. Application to very dry sandy soils followed by soaking rainfall may cause turf damage. Turf damage may also be increased in highly alkaline soils (soil pH >8.5 as determined by soil in water suspension). Avoid excessive mechanical disruptions such as aerification and verticutting within one week prior to or after application.

**RESISTANT WEEDS WARNING**

<b>GROUP</b>	<b>B</b>	<b>HERBICIDE</b>
--------------	----------	------------------

TRIBUTE Selective Turf Herbicide is a member of the sulfonylurea group of herbicides and has the inhibitor of ALS mode of action. For weed resistance management TRIBUTE is a Group B herbicide. Some naturally-occurring weed biotypes resistant to TRIBUTE, and other Group B herbicides, may exist through normal genetic variability in any weed population. These resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by TRIBUTE or other Group B herbicides. Do not rely exclusively on TRIBUTE for weed control. Use as part of an integrated weed management program involving herbicides with other modes of action and non-chemical methods of control. Since occurrence of resistant weeds is difficult to detect prior to use Bayer CropScience Pty. Ltd. accepts no liability for any losses that may result from the failure of TRIBUTE to control resistant weeds.

**MIXING**

Ensure that the spray tank is completely clean prior to mixing. Half fill the spray tank with water, then with agitators in motion, add the correct amount of TRIBUTE directly into the spray tank. Complete filling the tank with agitators in motion. Agitation must continue before and during spraying. If pH of water carrier is less than 5.5, use a buffer solution to raise pH to meet 7.0. **DO NOT** mix TRIBUTE with acid-forming compounds in the spray vat. **DO NOT** leave spray mix standing in the vat overnight.

**APPLICATION**

Ensure spraying equipment is properly calibrated before use. Ensure that complete and even spray coverage of all weeds is achieved. DO NOT overlap sprayed areas.

**Broadcast Application:**

Aim to apply a thorough and even coverage of spray to the target weed. Dense stands of weeds should be thoroughly wetted with spray. Incomplete coverage may result in poor control. Equipment set-up should be such that adequate coverage, penetration and volume of spray liquid can be achieved while the potential for off-target movement is minimised.

Application should be at the recommended rate in sufficient water to give thorough coverage of weeds. Apply in a minimum of 200 L/ha through nozzles that will deliver a MEDIUM spray droplet as defined by ASAE S572 Standard or BCPC Guideline. Use higher spray volumes (up to 500 L/ha) for dense weed populations or under adverse growing conditions.

The product should dry on the leaf surface prior to allowing traffic on the treated area. Risk to sensitive grasses such as bentgrass, ryegrass, and *Poa trivialis* adjacent to TRIBUTE applications is decreased when the soil is at less than field capacity. **DO NOT** apply if heavy rain has been forecast within 48 hours. **DO NOT** apply to waterlogged soil. TRIBUTE is foliar absorbed; irrigation is not recommended immediately after application. However, for the next two mornings after application, if dew is present, irrigate lightly (3-5 mm) prior to allowing traffic each day in the area. Avoid water run-off during these irrigation cycles. **DO NOT** irrigate to the point of run-off within 7 days of application.

**SPRAYER CLEAN UP**

The sprayer must be thoroughly decontaminated before being used again to spray susceptible plants or turf. Ensure that the following operation is carried out in an area that is clear of waterways, desirable vegetation and tree roots, and preferably in an area where drainings can be contained.

1. Drain sprayer completely and wash out tank, boom and hoses with clean water.
2. Drain again.
3. Fill the tank with clean water and add 300 mL of chlorine bleach (containing 4% chlorine) per 100 L of water with agitation running.
4. Flush some bleach solution through booms and hoses and allow remainder to agitate in tank for 10 minutes.
5. Remove nozzles and filters and leave to soak in a bleach solution of 500 mL per 10 L of water while tank cleaning is in progress.
6. Briefly run the pump at periodic intervals to refresh chlorine solution in spray lines.
7. Drain tank and repeat the procedure of flushing with bleach solution.
8. Flush the tank, boom and hoses with clean water.

**COMPATIBILITY**

No information is available that indicates physical compatibility with products other than those detailed in the Direction for Use table. Tank-mixing other products with TRIBUTE Selective Turf Herbicide cannot therefore be recommended. As formulations of other manufacturers' products are beyond the control of Bayer, all mixtures should be tested prior to mixing commercial quantities. Note that organophosphate insecticides should not be tank-mixed or applied within 7 days prior or 3 days after application as there may be increased potential for turf injury.

**PRECAUTIONS****RE-ENTRY PERIOD**

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to neck and wrist (or equivalent clothing), chemical resistant gloves and footwear. Clothing must be laundered after each days use.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. **DO NOT** use clippings from treated areas for mulch around vegetables or fruit trees.

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

TRIBUTE Selective Turf Herbicide may kill or harm non-target beneficial invertebrates where IPM is practised. Very toxic to aquatic life. **DO NOT** contaminate streams, rivers or waterways with this product or used containers. **DO NOT** apply if heavy rain has been forecast within 48 hours. **DO NOT** graze treated turf or feed turf clippings from any treated area to poultry or livestock.

Bayer Environmental Science  
A Business Operation of Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
Hawthorn East, Vic, 3123

Phone: (03) 9248 6888  
Technical enquiries: 1800 804 479  
Website: [www.bayeres.com.au](http://www.bayeres.com.au)