

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Pennmag™

Turf herbicide

syngenta®

ACTIVE CONSTITUENT: 960 g/L S-METOLACHLOR

GROUP **K** HERBICIDE

For pre-emergent control of weeds in turf as per the
Directions for Use

Syngenta Australia Pty Ltd

Level 1, 2-4 Lyonpark Road, Macquarie Park NSW 2113

**In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111
(24 hours)**

APVMA Approval No: 69865/102444

TM

DIRECTIONS FOR USE

Restrictions

- DO NOT apply by aircraft
- DO NOT apply using irrigation equipment
- DO NOT apply using spraying equipment carried on the back of the user
- DO NOT apply to turf under heat or moisture stress
- DO NOT apply to waterlogged soils
- DO NOT irrigate to the point of runoff for at least 3 days after application
- DO NOT apply if heavy rain has been forecast within 48 hours
- DO NOT apply to golf putting greens or bowling greens
- DO NOT apply to newly laid turf sods. Wait until turf has established
- DO NOT apply to newly seeded or recently over seeded turf. Wait until turf has established

Situation	Weed	Rate	Critical Comments
Turf	African Love Grass (<i>Eragrostis curvula</i>), Annual Ryegrass (<i>Lolium rigidum</i>), Barnyard Grass (<i>Echinochloa</i> spp), Burr Medic (<i>Medicago polymorpha</i>), Carrot Weed (<i>Cotula australis</i>), Crowsfoot Grass (<i>Eleusine indica</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Pigeon Grass (<i>Setaria</i> spp), Summer Grass (<i>Digitaria sanguinalis</i>) Winter Grass (<i>Poa annua</i>) Suppression of Pigweed (<i>Portulaca oleracea</i>)	2 L/ha or 20 mL/100 m ²	Apply prior to weed emergence for residual control of up to 8 weeks. Follow application with at least 3 mm irrigation. Sporting fields: Apply once per year in late winter to early spring. Turf production: Apply a single application to the soil immediately post-harvest. Refer to Application section for detailed information.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD: DO NOT GRAZE TREATED TURF/LAWN OR FEED TURF/LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK

GENERAL INSTRUCTIONS

PENNMAG is a pre-emergent herbicide for the control of various annual grasses and broadleaf weeds. PENNMAG is safe to use on established and recovering turf. The product acts mainly through shoot (coleoptile) absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the top 3 to 4 cm of soil, where it must be present at the time of weed germination. If applying PENNMAG to warm season turf during the transition from winter to spring, avoid applications when soil temperatures remain low. Turf recovery will be greater if applied when soils have already commenced warming.

Mixing

Add PENNMAG directly to a spray tank containing 2/3 of the required spray volume. Add the rest of the water and ensure the mix is thoroughly agitated before application. Maintain agitation during application.

Application

- DO NOT apply by aircraft.
- DO NOT apply using irrigation equipment.
- DO NOT apply using spraying equipment carried on the back of the user.
- Apply prior to weed emergence. Total application volume should be minimum 750 L/ha. Use coarse droplets (eg Air Induction flat fan 08 nozzles). Spray nozzles should be uniformly spaced and of the same size, and should provide accurate and uniform application. To ensure accuracy, calibrate sprayer at the beginning of the season before use and recalibrate frequently. Observe sprayer nozzles frequently during the spraying operation to ensure that the spray pattern is uniform. Avoid overlapping of spray runs. Ensure that boom height does not exceed 50 cm above the leaf blades of the turf.
- Fallow Soil:** When applying to fallow soil follow application with irrigation of at least 3 mm within 24 to 48 hours.
- Established Turf:** When applying to established turf (full or recovering canopy) **immediately** follow application with at least 3 mm irrigation.

Compatibility

As formulations of other manufacturers products are beyond the control of Syngenta, all mixtures should be tested before addition to a spray tank and before application to turf to establish safety.

Resistant Weeds Warning

GROUP K HERBICIDE

PENN MAG Turf Herbicide is a member of the amide group of herbicides. The product has the herbicides with diverse sites of action mode of action. For weed resistance management the product is a Group K herbicide. Some naturally occurring weed biotypes resistant to the product and other Group K herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group K herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Syngenta representative.

Integrated Pest Management

The possible effects of PENN MAG on integrated pest management (IPM) strategies in the turf industry have not been studied at the proposed rates.

PRECAUTION

Re-entry Period

General Public: DO NOT allow public re-entry into treated areas until the spray has dried.

Ground Staff / Professional Operators: DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If no facility is available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. DO NOT inhale spray mist. Will damage the eyes. Avoid contact with eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing.

When opening the container, preparing and using the prepared spray wear:

- goggles
- a disposable mist face mask covering mouth and nose

If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 067 108, or visit our website at www.greencast.com.au or www.syngenta.com.au

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

