

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Bronco<sup>®</sup> MA Herbicide

ACTIVE CONSTITUENTS: 200 g/L BROMOXYNIL  
present as the N-OCTANOYL ESTER  
200 g/L MCPA present as the ISO-OCTYL ESTER  
SOLVENT: 340 g/L LIQUID HYDROCARBON

GROUP **CI** HERBICIDE

For control of broadleaf weeds in cereals, linseed,  
grass pastures and turf as specified in the  
Directions for Use table.



ADAMA  
ESSENTIALS

adama.com

CONTENTS: 5 L, 20 L, 200 L, 110 L, 1000 L

### DIRECTIONS FOR USE

**Restrains:** DO NOT apply if crops, pastures or weeds are stressed due to dry or excessively moist conditions.  
DO NOT apply to diseased or frost-affected crops or if frosts are imminent.  
DO NOT apply if rain is expected within three hours.

SITUATION & CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Barley, Cereal Rye, Oats and Triticale;	Amsinckia ( <i>Amsinckia</i> spp.), Capeweed ( <i>Arctotheca Calendula</i> ), Chamomile ( <i>Matricaria matricarioides</i> ), Charlock ( <i>Sinapis arvensis</i> ), Corn Gromwell (Ironweed, Sheepweed), ( <i>Buglossoides arvensis</i> ), Fat Hen ( <i>Chenopodium album</i> ), Field Madder ( <i>Sherardia arvensis</i> ), Hexham Scent (King Island Melilot) ( <i>Mellilotus indicus</i> ), Horned Poppy ( <i>Glaucium flavum</i> ), Indian Hedge Mustard ( <i>Sisymbrium orientale</i> ), Lesser Swinecress ( <i>Coronopus didymus</i> ), Mexican Poppy ( <i>Argemone mexicana</i> ), Mintweed ( <i>Salvia reflexa</i> ), Paterson's Curse (Salvation Jane) ( <i>Echium plantagineum</i> ), Rough Poppy ( <i>Papaver hybridum</i> ), Saffron Thistle ( <i>Carthamus lanatus</i> ), Shepherd's Purse ( <i>Capsella bursa-pastoris</i> ), Slender Thistle ( <i>Corduus tenuiflorus</i> , <i>C. pycnocephalus</i> ), Tree Hogweed ( <i>Polygonum patulum</i> ), Turnip Weed (Giant Mustard) ( <i>Rapistrum rugosum</i> ), Wild Radish ( <i>Raphanus raphanistrum</i> ), Wild Turnip ( <i>Brassica tournefortii</i> )	Plants up to the 6 leaf stage but not more than 50 mm in diameter	WA only	750 mL	<b>CROP STAGE:</b> <b>Wheat, Barley, Cereal Rye, Oats and Triticale:</b> 3 leaf to fully tillered stage (Zadok Scale Z13-30) for rates of 750 mL to 1.4 litres per hectare; 5 leaf to fully tillered stage (Z15-30) for rates of 2 litres per hectare (NSW, Vic, SA, WA, Tas only); or early tillering to fully tillered stage (Z21-30) (Qld only).  <b>Linseed:</b> 5 to 15 cm high. DO NOT apply to linseed by aircraft.  # If spraying Black Bindweed when wheat, barley, cereal rye, oats, triticale or linseed is more than 30 cm high, increase spray volume to 200 litres per hectare.  For optimal control, apply only when the weeds are actively growing. In the early stages following spraying, temporary wilting of crop and pasture may occur. Crop scorch may occur, particularly on oats, if treatment is delayed to later stages of growth. DO NOT apply to crops or pastures undersown with lucerne, clover or medics unless some damage is acceptable. Avoid application during temperatures above 20° C.
		Plants up to the 8 leaf stage but not more than 75 mm in diameter	Tas, SA, NT only	2.0 L	
		WA only	1.4 L		
Linseed;					
Grass Pastures					

SITUATION & CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS	
Wheat, Barley, Cereal Rye, Oats and Triticale; Linseed; Grass Pastures – continued	Black Bindweed ( <i>Polygonum convolvulus</i> ) (Climbing Buckwheat), Common Peppergrass ( <i>Lepidium hyssopifolium</i> ), Mustards ( <i>Sisymbrium</i> spp.), Purple Calandrinia ( <i>Calandrinia menzieslii</i> )	Plants up to the 4 leaf stage but not more than 3.5 cm in diameter	WA only	1 L	-continued from previous page	
		Plants up to the 6 leaf stage but not more than 5 cm in diameter	All States	1.4 L		
		Plants up to the 8 leaf stage but not more than 7.5 cm in diameter		2.0 L		
	Fumitory ( <i>Fumaria</i> spp.), Three-cornered Jack (Doublegee, Spiny Emex) ( <i>Emex Australis</i> )	Plants up to the 3 leaf stage but not more than 5cm in diameter	Qld, NSW, Vic, Tas, SA only	1.4 L		
		Plants up to the 4 leaf stage but not more than 6 cm in diameter	WA only	750 mL		
		Plants up to the 5 leaf stage but not more than 7.5 cm in diameter	Qld, NSW, ACT, Vic, Tas, SA, NT only	2.0 L		
		Plants up to the 6 leaf stage but not more than 8.5 m in diameter	WA only	1.4 L		
	Variegated Thistle ( <i>Silybum marianum</i> ), Wireweed (Hogweed) ## ( <i>Polygonum aviculare</i> )	Plants up to the 4 leaf stage but not more than 3.5 cm in diameter	WA Only	750 mL		## Where Wireweed occurs in red soils of low fertility it has been found to be less susceptible. In these circumstances use 2 litres per hectare.
		Plants up to the 6 leaf stage but not more than 5 cm in diameter		Qld, NSW, ACT, Vic, Tas, SA, NT only		
		Mountain Sorrel ( <i>Oxalis acetosella</i> ), Three-horned Bedstraw ( <i>Galium tricornutum</i> )	Plants up to the 6 leaf stage	WA only		
	Plants up to the 8 leaf stage		750 mL			
	Cleavers ( <i>Galium aparine</i> )	2-4 stem stage and 1-3 whorls of leaves per stem		1.4 L		
	Turf: Established Couch (Including Queensland Blue), Kikuyu, Paspalum, Buffalo, Fescue, Bent and Rye lawns	Bindy-eye ( <i>Calotis hispidula</i> ), Capeweed ( <i>Arctotheca calendula</i> ), Catsear ( <i>Hypochoeris radicata</i> ), Common Plantain ( <i>Plantago</i> spp.), Creeping Buttercup ( <i>Ranunculus repens</i> ), Cudweed ( <i>Gnaphalium</i> spp.), Dandelion ( <i>Taraxacum officinale</i> ), Fleabane ( <i>Conyza</i> spp.), Jo-Jo (Onehunga) ( <i>Soliva pterosperma</i> )	-	All States		
Clover ( <i>Trifolium</i> spp.), Oxalis ( <i>Oxalis</i> spp.)			Qld, NSW, Vic, SA, WA only			

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**CERIALS, PASTURES:**

**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.**

**CROP HARVEST:**

**NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

### RESISTANT WEEDS WARNING

BRONCO® MA Herbicide contains members of the Nitrite and Phenoxy groups of herbicides. BRONCO MA has the photosystem II inhibitor and plant cell growth disruptor mode of action. For weed resistance management BRONCO MA contains Group C and I herbicides.

**GROUP CI HERBICIDE**

Some naturally-occurring weed biotypes resistant to one or more of the modes of action of BRONCO MA and other Group C and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by BRONCO MA or other Group C and I herbicides to whose mode of action they are resistant.

Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia Pty. Ltd. accepts no liability for any losses that may result from the failure of BRONCO MA to control resistant weeds. However resistance should not be assumed without first viewing the method of application, the coverage and the timing of application.

### MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

### APPLICATION

**BOOM SPRAYER:** Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. Wash the spray equipment thoroughly after use.

**CEREALS:** Apply 50-200 litres of water per hectare. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential.

**LINSEED:** Apply a minimum of 220 litres of water per hectare.

**TURF:** Apply a minimum of 500 litres of water per hectare.

**AERIAL APPLICATION:** Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microneaire equipment may give variable results.

### COMPATIBILITY

This product is physically compatible in the spray tank with one of the following herbicides and insecticides in wheat and barley only: metsulfuron methyl, chlorpyrifos, dicamba, dimethoate, endosulfan, aralkoxydim, diclofop-methyl, MCPA and omethoate.

When mixed with metsulfuron methyl or dicamba, DO NOT apply to cereals undersown with lucerne.

When mixed with diclofop-methyl, DO NOT exceed 1.4 litres per hectare.

As formulations of other manufacturers' products are beyond the control of Adama Australia Pty. Ltd. all mixtures should be tested prior to mixing commercial quantities.

### PROTECTION OF LIVESTOCK

#### *Grazing Precaution*

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where Capeweed, Paterson's Curse and Variegated Thistles predominate in the pasture. Avoid grazing with young or breeding stock.

DO NOT graze horses or pigs on Paterson's Curse. If in doubt, contact your nearest Department of Agriculture.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply by misting machines.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

### PROTECTION WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL (5 L, 20 L, 200 L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**For refillable containers,** empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### Envirodrum Micro Matic Valve (110 L)

Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of Adama Australia Pty. Ltd.

### 1000 L

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Adama Australia Pty. Ltd. should be advised immediately. This minibulk container is reusable and remains the property of Adama Australia Pty. Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Adama Australia Pty. Ltd. for cleaning, relabelling and refilling.

### SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. Attacks the eyes. Will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water.

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for BRONCO MA is available from Adama Australia on request. Call Customer Service on (02) 9431 7800.

**CONDITIONS OF SALE:** The use of BRONCO® MA Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia Pty. Ltd. regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia Pty. Ltd. accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered trademark of an ADAMA Agricultural Solutions Ltd Company

### NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

**Adama Australia Pty. Ltd.** ABN 55 050 328 973  
Suite 1, Level 4, Building B,  
207 Pacific Highway  
St Leonards NSW 2065 Australia  
Tel: (02) 9431 7800 Fax: (02) 9431 7700

APVMA Number: 47667/101042

Batch No.

Date of Manufacture