

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Revere[®] 500

Herbicide

ACTIVE CONSTITUENT: **500 g/L PROPYZAMIDE**

SOLVENT: 42 g/L ETHYLENE GLYCOL

GROUP **D** HERBICIDE

For selective control of certain grasses and broadleaf weeds in Lettuce, Sports Turf, Home Lawns and Legume Seed Crops and Pastures as per the DIRECTIONS FOR USE table

Formulation type
Suspension
Concentrate

SC



ADAMA

adama.com

CONTENTS: 5 L - 1000 L

DIRECTIONS FOR USE

| CROPS | WEEDS CONTROLLED | STATE | RATE L/ha | CRITICAL COMMENTS | |
|------------------------------|---|---------------|---------------------------------|---|---|
| Lettuce (Direct Sown) | Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, | All States | 4.5 in 330 - 550 L | Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours. | |
| Transplanted | Winter Grass, Ryegrass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherds Purse, Nettles | | | Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 - 4 cm of soil should be applied within 24 hours. | |
| | | Vic only | 2.3 in 300 - 550 L | Apply on light sandy soils, low in organic matter. | |
| Turf & Lawns Common Couch | Winter Grass | All States | 1.2 (12 mL/100 m ²) | Rate/ Bowling Green 37 x 37 m 170 mL | Apply during the Autumn and Winter, if established turf is infested with seedling Winter Grass. Ensure a uniform coverage. Irrigate immediately after application with 12- 25 mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Queensland Blue Couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on Bluegrass, Ryegrass, Fescue or Bent Grass Lawns. |
| Queensland Blue Couch | | Qld only | 0.6 (6 mL/100 m ²) | 85 mL | |
| Buffalo Grass | | WA only | 1.2 (12 mL/100 m ²) | 170 mL | |

ADAMA

| CROPS | WEEDS CONTROLLED | STATE | RATE L/ha | CRITICAL COMMENTS |
|--|--|----------------------------|-----------|---|
| Lucerne, Clover, Medics etc., grown for Forage, Hay or Seed production | Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue | NSW, ACT, SA, WA, Tas only | 1 - 1.5 | SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post or pre-emergent states. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended. |
| | Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed | | 1.5 - 2 | |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil. Mixing prior to pouring, shake container vigorously, then add the required quantity of Revere 500 Herbicide to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

RESISTANCE WARNING

REVERE[®] 500 Herbicide is a member of the benzamide group of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management this is a Group D Herbicide. Some naturally occurring weed biotypes resistant to REVERE[®] 500 Herbicide and Group D Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by REVERE[®] 500 Herbicide or Group D Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of REVERE[®] 500 Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary industries Department or an Adama representative.



PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

DO NOT graze sheep, on treated areas or feed to poultry or pigs.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for REVERE[®] Herbicide is available from adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of REVERE[®] Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

[®] Registered trademark of an Adama Group Company
12/2017 9559