

**CAUTION****KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING****Merit<sup>®</sup>****Turf and Ornamental Insecticide****ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID****GROUP 4A INSECTICIDE**

\*

For the control of a range of insect pests in turf and ornamental situations as per the directions for use  
IMPORTANT: Read the attached booklet before use.

\*1L, 2.5L and 5L pack sizes

**STORAGE AND DISPOSAL:** Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. DO NOT burn empty containers or product.

**SAFETY DIRECTIONS:**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID:**

If poisoning occurs contact a doctor or Poisons Information Centre (ph. 131126).

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which may be found at [www.bayeres.com.au](http://www.bayeres.com.au).

**Exclusion of Liability**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

**FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY:  
PHONE 1800 033 111**

Bayer Environmental Science  
Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn, Vic. 3123  
Phone: (03) 9248 6888  
Technical enquiries: 1800 804 479  
Website: [www.bayeres.com.au](http://www.bayeres.com.au)

MERIT<sup>®</sup> is a Registered Trademark of Bayer.

APVMA Approval No.: 59696 / 50036

BATCH NO:

DATE OF MANUFACTURE:

(\**drumMUSTER* logo on containers 2.5 L and larger)



**Directions for use**

<b>SITUATION</b>	<b>PEST</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m <sup>2</sup>	Apply at peak egg hatch, which is mid spring to mid summer depending on species.
	Larvae of billbug	Spray with at least 400 L water per hectare to ensure even coverage. Preferably spray on to wet or dewy grass.  Irrigate with 5 to 12 mm of water commencing within one to 24 hours of application.	Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.
Roses	Aphids	25 mL/100 L	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids Azalea lace bug Bronze orange bug Harlequin bug Citrus mealybug Greenhouse thrips Fullers rose weevil		
	Hibiscus flower beetle	50 mL/100 L	Spray buds and flowers as needed.
	Longtailed mealybug	50 mL/100 L + surfactant	Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	25 mL/100 L	Spray at first sign and then a week later.
	Soft scales		Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid		Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus trees	Flatid ( <i>Jamella australiae</i> )	<b>SpotSpray</b> 875 mL/100 L of water  <b>Stem injection</b> 1.75 L/1 L of water	<b>Spot spray:</b> Spray 100 mL of mixture directly into the leafy throat of each head. <b>Stem injection:</b> Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. DO NOT re-apply in the same holes. Uptake of MERIT, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

Situation	Pest	Rate	Critical Comments
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height	Use as a soil drench application. Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400-500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7-10 days after treatment. Treat at least 6-10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant	Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. DO NOT dilute to the point where mix runs out the bottom of pots.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot	Use as a soil drench for pots up to 20 L capacity.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water	Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.
Roses	Aphids	3.5 mL/2 L water/plant	Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of MERIT to the 2 L of water.
For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the MERIT mixture, then water it in well immediately after application.			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**GENERAL INSTRUCTIONS:**

**INSECTICIDE RESISTANCE WARNING:**

<b>GROUP</b>	<b>4A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For insecticide resistance management MERIT Turf and Ornamental Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to MERIT and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if MERIT or other Group 4A insecticides are used repeatedly. The effectiveness of MERIT on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of this product to control resistant insects.

MERIT Turf and Ornamental Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer Environmental Science representative. Insecticide Resistance Management Strategy for aphids other than green peach aphid, whitefly and melon thrips: DO NOT apply MERIT in consecutive sprays. Rotate with registered insecticides from other groups.

Confined environments such as glasshouses: Annuals: DO NOT apply more than one spray of MERIT to any one crop. Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

**PROTECTION OF LIVESTOCK:** Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT graze treated turf or feed turf clippings from any treated area to poultry or livestock.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:** DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**Note on Ornamentals**

MERIT Turf and Ornamental Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

**Mixing**

Prior to pouring, shake container vigorously then add the required quantity of MERIT Turf and Ornamental Insecticide to water in the spray vat while stirring or with agitators in motion. MERIT should not be mixed with other pesticides or fertilisers.

MERIT<sup>®</sup> is a Registered Trademark of Bayer.

Bayer Environmental Science  
Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn, Vic. 3123  
Phone: (03) 9248 6888  
Technical enquiries: 1800 804 479  
Website: [www.bayeres.com.au](http://www.bayeres.com.au)

APVMA Approval No.: 59696 / 50036