

MAIN PANEL

---

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Bayfidan<sup>®</sup>

## TURF FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L TRIADIMENOL  
SOLVENT: 602 g/L N-METHYL-2-PYRROLIDONE

GROUP	<b>C</b>	FUNGICIDE
-------	----------	-----------

1 L

For the systemic control of dollar spot and other diseases of turf as specified in the directions for use table

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE**

(label code)

FRONT PAGE OF BOOKLET

---

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Bayfidan<sup>®</sup>

## TURF FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L TRIADIMENOL  
SOLVENT: 602 g/L N-METHYL-2-PYRROLIDONE

GROUP	<b>C</b>	FUNGICIDE
-------	----------	-----------

1 L

For the systemic control of dollar spot and other diseases of turf as specified in the directions for use table

**IMPORTANT: READ THIS BOOKLET BEFORE USE**

(label code)

BOOKLET

---

**Protection of Crops, Native and Non-target Plants**

Do not apply earlier than 5 weeks after sowing.

**Protection of Wildlife, Fish, Crustaceans and Environment**

Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**Storage and Disposal**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of at local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**Safety Directions**

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

**First Aid**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126).

**Material Safety Data Sheet**

Additional information is listed in the Material Safety Data Sheet

**Exclusion of Liability**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Bayfidan® and Baythroid® are Registered Trademarks of Bayer.

APVMA Approval No.: 30676/1 L/1005

Bayer Environmental Science  
Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn Vic. 3123  
Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)

BOOKLET contd

**DIRECTIONS FOR USE**

Crop	Disease	Rate	Critical Comments
Turf	Dollar spot (Not Tasmania, NT)	Preventive 30 mL/100 m <sup>2</sup>	Apply thoroughly on a monthly program starting immediately the disease appears.
		Curative 60 mL/100 m <sup>2</sup>	Apply thoroughly under conditions of heavy disease pressure; then treat as necessary.
	Black Helminthosporium (QLD NSW TAS ONLY)	60 mL/100 m <sup>2</sup>	Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Apply during period of turf growth on a monthly program.
	Take-all patch (Not SA, NT)	60 mL/100 m <sup>2</sup>	Preventive only after soil fumigation with methyl bromide. Apply thoroughly, using high volume equipment, 6-8 weeks after seeding. Repeat applications (every 8 weeks in the first two years) for continued protection.
	Winter Fusarium (NSW VIC TAS ONLY)	60 mL/100 m <sup>2</sup>	Apply thoroughly on a monthly program from end of March until danger of infection has passed.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS****Fungicide Resistance Warning**

GROUP	<b>C</b>	FUNGICIDE
-------	----------	-----------

Bayfidan Turf Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide. Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group C fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

**Mixing:** Add the required quantity of Bayfidan to water in the spray vat while stirring or with agitators in motion.

Bayfidan is a broad spectrum, systemic, foliar fungicide that is quickly taken up by plants and translocated within the plant. Bayfidan provides protection on a monthly spray program and acts curatively after infection has taken place. Because of its systemic activity, Bayfidan is less affected by rainfall or irrigation than contact type fungicides.

**Compatibility**

Bayfidan is compatible with Baythroid® Turf Insecticide. The compatibilities given are based on world wide experience and are considered correct. However as changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, we accept no responsibility for the behaviour of such mixtures.

BACK PANEL ATTACHED TO CONTAINER

---

### Storage and disposal

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of at local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### Safety Directions

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

### First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 131126).

### Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet

### Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

**IMPORTANT: READ ATTACHED BOOKLET BEFORE USE**

APVMA Approval No.: 30676/1 L/1005

Bayfidan® and Baythroid® are Registered Trademarks of Bayer.

Bayer Environmental Science  
Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn Vic 3123  
Phone: 03 9248-6888  
Fax: 03 9248-6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE **1800 033 111**

\*Batch Number:

\*Date of Manufacture:

Bayer Environmental Science  
**A Business Group of Bayer CropScience**